

USER GUIDE

SOMRITH

Organic sapropel and peat-based biostimulant

SOMRITH is a powerful biostimulant containing 50% sapropel i.e. carbon-rich prehistoric aquatic organic matter. This 100% natural, mineral-rich composition of SOMRITH increases your soil organic matter and stimulates microbiological activity. Plants treated with SOMRITH show exceptional root and shoot growth and are more resistant to plant diseases—thanks to its formula containing humic and fulvic acid, chelated micronutrients, and amino acids derived from high-grade organic Russian peat.*

SOMRITH unlocks the power of sapropel to give you an abundant harvest.

*Upon following recommended dosage.

FEATURES & BENEFITS:

- ✓ Contains chemical-free, completely organic ingredients
- ✓ Contains chelated micronutrients for better nutrient absorption
- ✓ Enhances plant growth and quality
- ✓ Improves plant metabolism and nutrient uptake
- ✓ Improves plant immunity
- ✓ Improves soil water holding capacity
- ✓ Improves soil organic matter and soil health
- ✓ Improves soil quality thus increasing soil productivity

CONTENTS mg/L

Based on test report from Innovative Scientific Research and Test Centre for general Application, Russia Report No. 53 dt 19 May 2020

Total organic carbon	24.20%
Ash	13.00%
Humic acid content to total carbon	28.00%
Fulvic acid content to total carbon	21.00%

COMPATIBILITY

SOMRITH is compatible with a wide range of pesticides and fungicides. If compatibility of SOMRITH with a given product is not known, it is suggested to test the same by combining them together at the recommended rate before preparing a mix. **Suitable for all crops and compatible with all other agricultural inputs.**

WARRANTY

No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risk and liabilities resulting from handling, use or application of this product.



KEEP OUT OF REACH OF CHILDREN

INSTRUCTIONS IF CONSUMED ACCIDENTALLY BY HUMANS OR ANIMALS



Drink plenty of water and fluids and consult your doctor if any adverse symptoms appear.

STORAGE AND DISPOSAL

Store in cool and dry place and away from food and feed stuff. Dispose empty container in an approved sanitary landfill by burying or recycling.

For more information on SOMRITH get in touch with our representatives.

Contact: +91-11-40190695

Email: customercare@farmics.com

Farm better, together.

DIRECTIONS FOR USE

SOMRITH can be used for foliar application, root treatment, and seed treatment.

I. Foliar spray, seed or root treatment for vegetables/plants

Vegetables/Plants	Qty to be used per acre	Time	Application method	Recommended ratio for dilution / preparing Working Solution in water/acre
Vegetables (beetroot, brinjal, cabbage, cucumber, flowers, garlic, gourds, onions, squash, tomato, pepper, potato etc.)	40 ml Seed treatment	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Leafy vegetables (carrot, fenugreek, lettuce, radish, spinach)	40 ml Seed treatment	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Berries (strawberry, raspberry)	40 ml Seed treatment	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Grain & industrial crops (wheat, rice, maize, barley, oats, sorghum, pearl millet (bajra) etc.)	40 ml Seed treatment	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Technical & forage crops (mustard, soybean, groundnut, sunflower, jute, forages)	40 ml Seed treatment	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Legumes (chickpea, pigeonpea, peas, beans, lentils, etc.)	40 ml	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Floriculture	40 ml	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine
Cotton	40 ml	Before sowing	Soaking of seeds for 6 hours	1 L water
	500 ml	30 days	Spray	100 L water and 25 L cow urine
	600 ml	60 days	Spray	120 L water and 30 L cow urine
	900 ml	90 days	Spray	180 L water and 45 L cow urine
Any other crop	40 ml	Before sowing	Soaking of seeds for 4 hours	1 L water
	500 ml	15 days	Root drenching	100 L water and 25 L cow urine
	600 ml	30 days	Spray	120 L water and 30 L cow urine
	900 ml	60 days	Spray	180 L water and 45 L cow urine

II. Replanting with shoots

Vegetables/Plants	Qty to be used	Time	Application method	Recommended ratio for preparing Working Solution in water/acre
Banana and other shoots	600 ml	At the time of planting	Soaking of plants / root drenching / shoots for 4 hours	15 L water
	800 ml	30 days	Spray / Through Drip Irrigation System	160 L water and 40 L cow urine
	800 ml	90 days	Through Drip Irrigation System	160 L water and 40 L cow urine
	600 ml	150 days	Through Drip Irrigation System	120 L water and 30 L cow urine
	600 ml	210 days	Through Drip Irrigation System	120 L water and 30 L cow urine

* For best results use the indicated dosage.